




THE *Animal* CLINIC of Kalispell

1408 Airport Road
Kalispell, MT 59901
(406) 755-6886
(406) 755-6898 FAX

What is a corneal ulcer?

The cornea is the transparent, shiny membrane which makes up the front of the eyeball. Think of it as a clear window.



How does a corneal ulcer occur?

There are several causes of corneal ulcers in dogs. The most common is trauma. An ulcer may result from blunt trauma, such as a dog rubbing its eye on the carpet, or due to a laceration, such as a cat scratch or a sharp object. Another common cause is chemical burn of the cornea. This may happen when irritating shampoo gets in the eye.

Less common causes of corneal ulcers include bacterial infections, viral infections, and other diseases. These may originate in the eye or may develop secondary to a disease elsewhere in the body.

How does a corneal ulcer affect my dog?

A corneal ulcer is very painful. Most dogs rub the affected eye with a paw or on the carpet in an attempt to relieve this intense pain. To protect the eye, they keep the lids tightly closed. Occasionally, there will be a discharge that collects in the corner of the eye or runs down the face.

How is a corneal ulcer diagnosed?

Superficial corneal abrasions are usually not visible without special tests. Corneal abrasions can be visualized with the use of special stains such as **fluorescein**. A drop of this stain is placed on the cornea. The dye will adhere to an area of ulceration and is easily seen using special ophthalmic lights. A fluorescein stain test is the most common eye test performed and may be the only test needed if the ulcer is acute and very superficial.

How is a corneal ulcer treated?

Corneal abrasions generally heal within three to five days. Medication is used to prevent bacterial infections (antibiotic ophthalmic drops or ointment. Antibiotic drops are only effective for a few minutes so they must be applied frequently; ointments last a bit longer but still require application every few hours. It is suggested that an antibiotic preparation be instilled in the eye every four to eight hours..



If a corneal ulcer is present, measures must be taken to protect the eye and to promote healing. Since dogs do not wear eye patches well, surgery may be required to protect the injury and allow for normal healing. In certain cases, it may be necessary to perform surgery to remove dead or poorly healing layers of corneal tissue, or to perform a corneal graft. Your veterinarian will choose the best surgery to optimize healing for your pet.

Are there any side-effects from the eye medications?

Occasionally a dog will be sensitive to an ophthalmic antibiotic. If your dog seems to be in more pain after the medication is used, discontinue it and contact your veterinarian immediately.

Since a corneal ulcer is painful, can I apply a topical anesthetic to the cornea?

A topical anesthetic is often used to numb the cornea so the diagnostic tests may be performed. However, these drugs often delay healing and should be used with discretion.

How do I know when to discontinue medication?

The best way to tell that the cornea has healed is to repeat the fluorescein stain test. This will be performed by your veterinarian five to seven days after treatment has begun.

There appear to be some red streaks near the ulcer. Is that normal?

The normal cornea has no blood vessels in it. However, when a corneal ulcer occurs, the body tries to heal by growing new blood vessels through a process called *neovascularization*. The new vessels begin at the sclera (the white part of the eye) and cross the cornea to the ulcer.

Neovascularization is a good response because it indicates healing. However, after the ulcer is healed, these vessels remain in the cornea. They are not painful, but they do obstruct vision. Therefore, it is desirable to minimize their size. This is done with the use of ophthalmic drops or ointment containing a corticosteroid, which may be used for a few days to several weeks, depending on how much neovascularization is present.

It is important that steroids are not used in the eye too soon because they will stop the healing process and may cause complications. Therefore, the fluorescein dye test should be performed before beginning this type of medication. If steroids are used and the eye becomes painful again, immediately discontinue the steroids and contact your veterinarian.